

# Deterministic Identity Testing of Read-Once Algebraic Branching Programs

Maurice Jansen\*

Youming Qiao\*

Jayalal Sarma M.N.\*

December 14, 2009

## Abstract

In this paper we study polynomial identity testing of sums of  $k$  read-once algebraic branching programs ( $\Sigma_k$ -RO-ABPs), generalizing the work of Shpilka and Volkovich [1, 2], who considered sums of  $k$  read-once formulas ( $\Sigma_k$ -RO-formulas). We show that  $\Sigma_k$ -RO-ABPs are strictly more powerful than  $\Sigma_k$ -RO-formulas, for any  $k \leq \lfloor n/2 \rfloor$ , where  $n$  is the number of variables. Nevertheless, as a starting observation, we show that the generator given in [2] for testing a single RO-formula also works against a single RO-ABP.

For the main technical part of this paper, we develop a property of polynomials called *alignment*. Using this property in conjunction with the *hardness of representation approach* of [1, 2], we obtain the following results for identity testing  $\Sigma_k$ -RO-ABPs, provided the underlying field has enough elements (more than  $kn^4$  suffices):

1. Given free access to the RO-ABPs in the sum, we get a deterministic algorithm that runs in time  $O(k^2 n^7 s) + n^{O(k)}$ , where  $s$  bounds the size of any largest RO-ABP given on the input. This implies we have a deterministic polynomial time algorithm for testing whether the sum of a constant number of RO-ABPs computes the zero polynomial.
2. Given black-box access to the RO-ABPs computing the *individual* polynomials in the sum, we get a deterministic algorithm that runs in time  $k^2 n^{O(\log n)} + n^{O(k)}$ .
3. Finally, given only black-box access to the polynomial computed by the sum of the  $k$  RO-ABPs, we obtain an  $n^{O(k+\log n)}$  time deterministic algorithm.

Items 1. and 3. above strengthen two main results of [2] (Theorems 2 and 3, respectively, for the case of non-preprocessed  $\Sigma_k$ -RO-formulas).

## 1 Introduction

In this paper we make contributions to the program of constructing increasingly more powerful pseudo-random generators useful against arithmetic circuits. As argued by Agrawal [3], this program is an approach towards resolving Valiant's Hypothesis, which states that the algebraic complexity classes VP and VNP are distinct.

Central to this program is the PIT problem: given an arithmetic circuit  $C$  with input variables  $x_1, x_2, \dots, x_n$  over a field  $\mathbb{F}$ , test if  $C(x_1, x_2, \dots, x_n)$  computes the zero polynomial in the ring

---

\*Institute for Theoretical Computer Science, Tsinghua University, Beijing, P.R. China. Email: maurice.julien.jansen@gmail.com, jimmyqiao86@gmail.com, jayalal@tsinghua.edu.cn. This work was supported in part by the National Natural Science Foundation of China Grant 60553001, and the National Basic Research Program of China Grant 2007CB807900, 2007CB807901.

$\mathbb{F}[x_1, x_2, \dots, x_n]$ . This is a well-studied algorithmic problem with a long history and a variety of connections and applications. See [4] for a recent survey. Efficient randomized algorithms were proposed independently by Schwartz [5] and Zippel [6]. Obtaining a deterministic algorithm for the problem seemed surprisingly elusive.

It was originally Kabanets and Impagliazzo [7] who showed the strong connection between derandomizing PIT and proving circuit lower bounds. They showed that giving a deterministic polynomial time (even subexponential time) identity testing algorithm means either that  $\text{NEXP} \not\subseteq \text{P}/\text{poly}$ , or that the permanent has no polynomial size arithmetic circuits. This was further strengthened in [3], where it was shown that giving a black-box derandomization of PIT implies that an explicit multilinear polynomial has no subexponential size arithmetic circuits.

Since the seminal work of [7], there has been a lot of attention and an impressive amount of progress in the area. Some of the special cases for which progress has been reported are: depth-2 arithmetic formulas [8, 9, 10], depth-3 and depth-4 arithmetic circuits with bounded top fan-in [11, 12, 13, 14, 15, 16], and non-commutative arithmetic formulas [17]. In a surprising result, Agrawal and Vinay [18] showed that the black-box derandomization of PIT for only depth-4 circuits is almost as hard as that for general arithmetic circuits.

Partly aimed at making progress towards an efficient deterministic PIT algorithm for multilinear formulas, Shpilka and Volkovich [1, 2] studied the arithmetic read-once formula model. An arithmetic read-once formula is given by a tree whose nodes are taken from  $\{+, \times\}$ , and whose leaves are variables or field constants, subject to the restriction that each variable  $x_i$  is allowed to appear at most once. In their work, efficient black-box deterministic PIT algorithms are given for  $\Sigma_k$ -RO-formulas, for “moderate”  $k$ .

We remark that due to a construction by Valiant [19], given a RO-formula  $F$  of size  $s$  computing  $f$ , one can express  $f$  as a “read-once” determinantal expression  $f = \det(M)$ , where  $M$  is a  $O(s)$ -dimensional matrix, whose entries are variables or field elements. In this, each variable  $x_i$  appears at most once in  $M$ . Identity testing read-once determinantal expressions, is an important special case of the PIT problem, as it is well-known that the bipartite perfect matching problem (BIPARTITE-PM) reduces to that form. Giving a black-box algorithm for testing such expressions has the potential of putting BIPARTITE-PM in NC, which is a prominent open problem in complexity theory regarding parallelizability [20, 21, 22, 23].

## 1.1 Results

We consider a generalization of the above mentioned RO-formulas, namely *read-once algebraic branching programs* (RO-ABP)<sup>1</sup>. An algebraic branching program (ABP) is a layered directed acyclic graph with two special vertices  $s$  and  $t$ . Each edge is assigned a weight, which is an element of  $X \cup \mathbb{F}$ , where  $X$  is a set of variables. For a path in the graph its weight is taken to be the product of the weight on its edges. The ABP itself computes a polynomial which is the sum of the weights of all paths from  $s$  to  $t$ . The ABP is said to be *read-once* if each variable appears on at most one edge. A polynomial  $f \in \mathbb{F}[X]$  is called a *RO-ABP-polynomial* if there exists a RO-ABP which computes  $f$ .

Due to [19], if  $f$  can be computed by a RO-formula of size  $s$ , then  $f$  can be computed by a RO-ABP of size  $O(s)$ . However, RO-ABPs are strictly more powerful than RO-formulas. Appendix A shows a RO-ABP computing  $g = x_1x_2 + x_2x_3 + \dots + x_{2n-1}x_{2n}$ . Example 3.12 in [1] shows that

---

<sup>1</sup>See Section 2 for a formal definition.

$g$  can not be computed by a RO-formula, if  $n \geq 2$ . We remark that the RO-ABP model is not universal, e.g. for  $n \geq 3$ ,  $\prod_{1 \leq i < j \leq n} x_i x_j$ , is not an RO-ABP-polynomial (See Appendix B). By [19], if  $f$  is computable by a RO-ABP of size  $s$ , then we can write  $f$  as a read-once determinantal expression  $f = \det(M(x))$ , where  $M$  is a matrix of dimension  $O(s)$ .

The results we will mention next make progress towards identity testing read-once determinantal expressions. This contributes to the program for separating VP and VNP mentioned in previous section (See e.g. [24] for a direct connection).

Our first result is to show that the Shpilka-Volkovich generator (SV-generator) used in [2] for identity testing RO-formulas also provides a test for RO-ABPs. This generator has also very recently been applied to identity testing multilinear depth 4 circuits with bounded top fan-in [16]. It is defined as follows:

Let  $A = \{a_1, a_2, \dots, a_n\} \subseteq \mathbb{F}$  be a set of size  $n$ . For every  $i \in [n]$ , let  $u_i(w)$  be the  $i$ th Lagrange interpolation polynomial on  $A$ . Then  $u_i(w)$  is a polynomial of degree  $n-1$  satisfying that  $u_i(a_j) = 1$  if  $j = i$  and 0 otherwise. For every  $i \in [n]$  and  $k \geq 1$ , define

$$G_k^i(y_1, y_2, \dots, y_k, z_1, z_2, \dots, z_k) = \sum_{j \in [k]} u_i(y_j) z_j.$$

and let  $G_k(y_1, y_2, \dots, y_k, z_1, z_2, \dots, z_k) : \mathbb{F}^{2k} \rightarrow \mathbb{F}^n$ , be defined by  $G_k = (G_k^1, G_k^2, \dots, G_k^n)$ . We refer to the polynomial mapping  $G_k$  as the  $k$ th-order SV-generator, or SV-generator for short. We have the following ‘‘Generator Lemma’’:

**Lemma 1.** *Let  $f \in \mathbb{F}[X]$  be a nonzero RO-ABP-polynomial with  $|\text{var}(f)| \leq 2^m$ , for some  $m \geq 0$ . Then  $f(G_{m+1}) \neq 0$ .*

To make further progress, we consider sums of  $k$  RO-ABPs. We give an explicit *hitting-set* of size  $n^{O(k+\log n)}$  for  $\Sigma_k$ -RO-ABPs. Namely we have the following theorem:

**Theorem 1.** *Let  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$  be a set of  $k$  RO-ABPs. Let  $f = \sum_{i \in [k]} f_i$ . Provided  $|\mathbb{F}| > kn^4$ , we have that  $f \equiv 0 \iff \forall a \in \mathcal{W}_k^n + \mathcal{A}_k, f(a) = 0$ , where  $\mathcal{W}_k^n = \{y \in \{0, 1\}^n \mid \text{wt}(y) \leq k\}$  and  $\mathcal{A}_k = G_m(V^{2m})$  for the  $m$ th-order SV-generator with  $m = \lceil \log n \rceil + 1$ , and  $V \subset \mathbb{F}$  is a arbitrary set of size  $kn^4 + 1$ .*

In the above for  $V, W \subseteq \mathbb{F}^n$ ,  $V + W$  denotes the set  $\{v + w : v \in V, w \in W\}$ . By Theorem 1, we obtain the following black-box PIT for  $\Sigma_k$ -RO-ABPs:

**Theorem 2.** *Let  $f = \sum_{i \in [k]} f_i$  be a sum of  $k$  RO-ABP-polynomials in  $n$  variables. Let  $\mathbb{F}$  be a field with  $|\mathbb{F}| > kn^4$ . Given black-box access to  $f$ , it can be decided deterministically in time  $n^{O(k+\log n)}$  whether  $f \equiv 0$ .*

This strengthens a main result of [2] (Theorem 3, for the non-preprocessed<sup>2</sup> case), which provides a deterministic  $n^{O(k+\log n)}$  time PIT algorithm for  $\Sigma_k$ -RO-formulas. Namely, we prove a strict separation between  $\Sigma_k$ -RO-formula and  $\Sigma_k$ -RO-ABP, for  $k \leq \lfloor n/2 \rfloor$ . We show that

**Theorem 3.**  $\prod_{i \in [2n], i \text{ is odd}} \prod_{j \in [2n], j \text{ is even}} x_i x_j$  can not be written as a sum of  $\lfloor n/2 \rfloor$  RO-formulas.

The polynomial of Theorem 3 can be computed by a *single* RO-ABP of size  $O(n^2)$  (see Section 3). In the non-black-box setting we will prove the following result:

---

<sup>2</sup>A generalization of our theorems to preprocessed  $\Sigma_k$ -RO-ABPs will not be pursued here.

**Theorem 4.** Let  $\{A_i\}_{i \in [k]}$  be a set of  $k$  RO-ABPs in  $n$  variables. Let  $\mathbb{F}$  be a field with  $|\mathbb{F}| > kn^2$ . Given  $\{A_i\}_{i \in [k]}$  on the input, it can be decided deterministically in time  $O(k^2 n^7 s) + n^{O(k)}$  whether  $\sum_{i \in [k]} f_i \equiv 0$ , where  $f_i$  is the RO-ABP-polynomial computed by  $A_i$ , for  $i \in [k]$ .

Since the construction in [19] can be computed efficiently, this strengthens Theorem 2 in [2], for the case of non-preprocessed  $\Sigma_k$ -RO-formulas.

Finally, if black-box access is granted to the individual  $f_i$ 's, which we call the *semi-black-box* setting, we obtain the following result:

**Theorem 5.** Let  $\{f_i\}_{i \in [k]}$  be a set of  $k$  RO-ABP-polynomials in  $n$  variables. Let  $\mathbb{F}$  be a field with  $|\mathbb{F}| > kn^2$ . Given black-box access to each individual  $f_i$ , it can be decided deterministically in time  $k^2 n^{O(\log n)} + n^{O(k)}$  whether  $\sum_{i \in [k]} f_i \equiv 0$ .

## 1.2 Techniques for $\Sigma_k$ -RO-ABP PIT

The results for  $\Sigma_k$ -RO-ABP PIT are obtained through the *hardness of representation* approach of [1, 2]. There the PIT algorithm is derived from a statement that  $x_1 x_2 \dots x_n$  cannot be expressed as a sum of  $k \leq n/3$  RO-formula computable polynomials  $\{f_i\}_{i \in [k]}$ , if the polynomials  $f_i$  satisfy some special property. We do not need to define this special property for the discussion here, except that we should name it:  $\bar{0}$ -justification.

Unfortunately, the property of  $\bar{0}$ -justification, does not work for the  $\Sigma_k$ -RO-ABP model. With some thought it can be seen that the monomial  $x_1 x_2 \dots x_n$  is expressible as the sum of three  $\bar{0}$ -justified RO-ABP-polynomials. Our main technical contribution is the development of a new “special property”, called *alignment*, for which a hardness of representation theorem can still be proved, but which also can be satisfied simultaneously for a collection of RO-ABP-polynomials by means of an efficiently computable coordinate shift.

With regards to the latter, consider  $f = f_1 + f_2 + \dots + f_k$ , where each  $f_i$  is a RO-ABP-polynomial. Observe that  $\forall v \in \mathbb{F}^n$ ,  $f \equiv 0 \iff f(x_1 + v_1, x_2 + v_2, \dots, x_n + v_n) \equiv 0$ . With some technical work, we will establish a *sufficient* condition for alignment. With it we show that we can compute a coordinate shift  $v$  such that all  $f_i(x + v)$  are aligned. Such a shift  $v$  is called a *simultaneous alignment*. In the case of having only black-box access to  $f$ , we will show we have a “small” set of candidates containing at least one simultaneous alignment. The PIT algorithms will follow from this.

The rest of this paper is organized as follows. Section 2 contains preliminaries. In Section 3 we compare  $\Sigma_k$ -RO-formulas and  $\Sigma_k$ -RO-ABPs. In Section 4 we prove Generator Lemma 1. In Section 5 we develop the tools regarding alignment. Then in Section 6 we show how to compute a simultaneous alignment. Section 7 contains the hardness of representation theorem for RO-ABPs. From these developments, we put the PIT algorithms together in Section 8.

## 2 Preliminaries

Let  $X = \{x_1, x_2, \dots, x_n\}$  be a set of variables and let  $\mathbb{F}$  be a field. Let  $\mathcal{W}_k^n = \{y \in \{0, 1\}^n \mid wt(y) \leq k\}$ , where  $wt(y)$  counts the number of ones in  $y$ .

**Definition 1.** (RO-ABPs) An algebraic branching program (ABP) is a 4-tuple  $A = (G, w, s, t)$ , where  $G = (V, E)$  is an edge-labeled directed acyclic graph for which the vertex set  $V$  can be parti-

tioned into levels  $L_0, L_1, \dots, L_d$ , where  $L_0 = s$  and  $L_d = t$ . Vertices  $s$  and  $t$  are called the source and sink of  $B$ , respectively. Edges may only go between consecutive levels  $L_i$  and  $L_{i+1}$ .

The label function  $w : E \rightarrow X \cup \mathbb{F}$  assigns variables or field constants to the edges of  $G$ . For a path  $p$  in  $G$ , we extend the weight function by  $w(p) = \prod_{e \in p} w(e)$ . Let  $P_{i,j}$  denote the collection of all directed paths  $p$  from  $i$  to  $j$  in  $G$ . The program  $A$  computes the polynomial  $\hat{A} := \sum_{p \in P_{s,t}} w(p)$ . The size of  $A$  is defined to be  $|V|$ .

An ABP is said to be *read-once* if  $|w^{-1}(x_i)| \leq 1$ , for each  $x_i \in X$ . That is, every variable is read at most once by the program. A polynomial  $f \in \mathbb{F}[X]$  is called a *RO-ABP-polynomial*, if there exists a RO-ABP which computes  $f$ . We use the following notation: for  $x_i$  present on arc  $(v, w)$  in a RO-ABP  $A$ :  $\text{begin}(x_i) = v$  and  $\text{end}(x_i) = w$ . We let  $\text{source}(A)$  and  $\text{sink}(A)$  stand for the source and sink of  $A$ . For any nodes  $v, w$  in  $A$ , we denote the subprogram with source  $v$  and sink  $w$  by  $A_{v,w}$ . A *layer* of a RO-ABP  $A$  is any subgraph induced by two consecutive levels  $L_i$  and  $L_{i+1}$  in  $A$ . We will assume RO-ABPs are in the form given by the following straightforwardly proven lemma:

**Lemma 2.** *If  $f \in \mathbb{F}[X]$  is a RO-ABP-polynomial, then  $f$  can be computed by a RO-ABP  $A$ , where every layer contains at most one variable-labeled edge.*

Let  $f$  be a polynomial in the ring  $\mathbb{F}[X]$ . For  $\alpha \in \mathbb{F}$ ,  $f|_{x_i=\alpha}$  denotes the polynomial  $f(x_1, x_2, \dots, x_{i-1}, \alpha, x_{i+1}, \dots, x_n)$ . Extending this to sets of variables, for a subset  $I \subseteq [n]$  and an assignment  $a \in \mathbb{F}^n$ ,  $f|_{x_I=a_I}$  is the polynomial resulting from setting the variable  $x_i$  to  $a_i$  in  $f$  for every  $i \in I$ . This is not to be confused with the following notation: for  $S \subseteq \mathbb{F}^n$ , we will write  $f|_S \equiv 0$  to denote that  $\forall a \in S, f(a) = 0$ .

The following two notions are taken from [2]. We say that a polynomial  $f$  *depends on a variable*  $x_i$  if there exists an  $a \in \mathbb{F}^n$  and  $b \in \mathbb{F}$ , such that  $f(a_1, a_2, a_{i-1}, a_i, a_{i+1}, \dots, a_n) \neq f(a_1, a_2, a_{i-1}, b, a_{i+1}, \dots, a_n)$ . The set of variables  $x_i$  that  $f$  depends on is denoted by  $\text{Var}(f)$ . For a polynomial  $f \in \mathbb{F}[X]$ , the *partial derivative with respect to  $x_i$* , denoted by  $\frac{\partial f}{\partial x_i}$ , is defined as  $f|_{x_i=1} - f|_{x_i=0}$ . We will freely use the properties listed for this notion in [2]. For example, a multilinear polynomial  $f$  depends on  $x_i$  if and only if  $\frac{\partial f}{\partial x_i} \neq 0$ . In addition,  $\frac{\partial f}{\partial x_i}$  does not depend on  $x_i$ . Partial derivatives commute, which we express by saying that  $\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}$ . Setting values to variables commutes with taking partial derivatives in the following way:  $\forall i \neq j, \frac{\partial f}{\partial x_i}|_{x_j=a} = \frac{\partial(f|_{x_j=a})}{\partial x_i}$ .

**Lemma 3.** *Let  $f \in \mathbb{F}[X]$  be a RO-ABP-polynomial, then  $\frac{\partial f}{\partial x_i}$  is a RO-ABP-polynomial.*

*Proof.* Let  $p = |\text{var}(f)|$ . In case  $p = 0$  it is trivial. Assume  $p > 0$ . If  $x_i \notin \text{var}(f)$ , then  $\frac{\partial f}{\partial x_i} \equiv 0$ , in which case the property trivially holds. Now suppose  $x_i \in \text{var}(f)$ . Hence  $x_i$  must appear somewhere in  $A$ . Say  $x_i$  is on the arc  $(v_1, w_1)$  from level  $L_j$  to  $L_{j+1}$ , where  $L_j = \{v_1, v_2, \dots, v_{m_1}\}$  and  $L_{j+1} = \{w_1, w_2, \dots, w_{m_2}\}$ , for certain  $j, m_1, m_2$ . We can write

$$f = \sum_{a \in [m_1]} \sum_{b \in [m_2]} f_{s,v_a} w(v_a, w_b) f_{w_b,t}, \quad (1)$$

where for any nodes  $p$  and  $q$  in  $A$ ,  $f_{p,q}$  is the polynomial computed by subprogram  $A_{p,q}$ . Then

$$\begin{aligned}
\frac{\partial f}{\partial x_i} &= f_{|x_i=1} - f_{|x_i=0} \\
&= \sum_{a \in [m_1]} \sum_{b \in [m_2]} f_{s,v_a} w(v_a, w_b)_{|x_i=1} f_{w_b,t} - \sum_{a \in [m_1]} \sum_{b \in [m_2]} f_{s,v_a} w(v_a, w_b)_{|x_i=0} f_{w_b,t} \\
&= \sum_{a \in [m_1]} \sum_{b \in [m_2]} f_{s,v_a} (w(v_a, w_b)_{|x_i=1} - w(v_a, w_b)_{|x_i=0}) f_{w_b,t} \\
&= f_{s,v_1} f_{w_1,t}.
\end{aligned}$$

Hence we obtain a valid RO-ABP computing  $\frac{\partial f}{\partial x_i}$  from  $A$  by setting the label of the wire  $(v_1, w_1)$  to 1, and removing all other wires between layers  $L_j$  and  $L_{j+1}$ .  $\square$

The proof of the above lemma provides the insight that a RO-ABP computing  $\frac{\partial f}{\partial x_i}$  can be obtained from a RO-ABP computing  $f$ , by setting  $x_i = 1$  and removing all other edges in the layer containing  $x_i$ . This fact will be used at several places in the paper. Finally, observe the following simple-but-useful factor-lemma:

**Lemma 4.** *If  $f \in \mathbb{F}[X]$  is a RO-ABP-polynomial such that  $f \not\equiv 0$  and  $f = g \cdot (\beta x_i - \alpha)$ , then  $g$  is a RO-ABP-polynomial.*

*Proof.* This follows from the fact that for every  $\gamma$  with  $\beta\gamma - \alpha \neq 0$ ,  $g = \frac{1}{\beta\gamma - \alpha} \cdot f_{|x_i=\gamma}$ .  $\square$

## 2.1 Combinatorial Nullstellensatz and a Lemma by Gauss

**Lemma 5** (Lemma 2.1 in [25]). *Let  $f \in \mathbb{F}[X]$  be a nonzero polynomial such that the degree of  $f$  in  $x_i$  is bounded by  $r_i$ , and let  $S_i \subseteq \mathbb{F}$  be of size at least  $r_i + 1$ , for all  $i \in [n]$ . Then there exists  $(s_1, s_2, \dots, s_n) \in S_1 \times S_2 \times \dots \times S_n$  with  $f(s_1, s_2, \dots, s_n) \neq 0$ .*

**Lemma 6.** (Gauss) *Let  $P \in \mathbb{F}[X, y]$  be a nonzero polynomial, and let  $g \in \mathbb{F}[X]$  be such that  $P|_{y=g(x)} \equiv 0$ . Then  $y - g(x)$  is an irreducible factor of  $P$  in the ring  $\mathbb{F}[X]$ .*

## 3 Separation of RO-ABP and $\Sigma_{\lfloor n/2 \rfloor}$ -RO-formulas

For  $n \geq 2$ , let  $f_n$  be defined as

$$f_n(x_1, x_2, \dots, x_{2n-1}, x_{2n}) = \prod_{i \in [2n], i \text{ is odd}} \prod_{j \in [2n], j \text{ is even}} x_i x_j.$$

**Proposition 1.**  *$f_n$  can be computed by an RO-ABP of size  $O(n^2)$ .*

*Proof.* The RO-ABP is shown in Figure 1. Note that between the  $(n+1)$ th level and the  $(n+2)$ th level there is an  $n$  by  $n$  complete bipartite graph.  $\square$

**Proposition 2.** *A polynomial  $p(x_1, x_2, \dots, x_n)$  that contains three terms of form  $\alpha x_i x_j + \beta x_j x_k + \gamma x_k x_l$ , where  $i, j, k, l \in [n]$  are pairwise different, and  $\alpha, \beta, \gamma \in \mathbb{F}$  are nonzero, can not be computed by a RO-formula, for  $n \geq 4$ .*




Figure 1: A RO-ABP computing  $f_n$ .

*Proof.* For the purpose of contradiction, suppose there is a RO-formula  $F$  computing  $p$ . Setting all  $x_m = 0$ , for  $m \in [n] \setminus \{i, j, k, l\}$ , would result in an RO-formula  $F'$  computing  $p'(x_i, x_j, x_k, x_l) = \alpha x_i x_j + \beta x_j x_k + \gamma x_k x_l + a x_i + b x_j + c x_k + d x_l + e$ . However,  $p'$  can not be computed by an RO-formula. One argues this in a similar manner as for  $x_1 x_2 + x_2 x_3 + x_3 x_4$  (See example 3.12 in [1]). □

Consider the complete bipartite graph  $G_n = (V_n, E_n)$  for  $f_n$ , called the graph associated with  $f_n$ , shown in Figure 2. Every edge represents a term in  $f_n$ . The term  $x_i x_j + x_j x_k + x_k x_l$  can be viewed as a length-3 path in  $G_n$ .

**Proposition 3.** *Let  $n \geq 2$ . In  $G_n$ , for an edge set  $S \subseteq E_n$  with  $|S| \geq 2n - 1$ ,  $S$  must contain a length-3 path.*

*Proof.* We just need to prove that for  $G_n$ , the maximum “length-3 path free” edge set is of size at most  $2(n - 1)$ . This is proved by induction on  $n$ . For  $n = 2$ , it is easy to see that it holds. Suppose for  $n < l$  the claim holds. Then for  $n = l$ , for any length-3 path free edge set  $S$ , consider the following two cases:

1. If there exists an edge  $e = (u, v) \in S$ , for which  $u$  or  $v$  has no other outgoing edges, let  $S' = S \setminus \{e\}$ .  $S'$  is a length-3 path free set in  $G_{l-1}$ . By induction,  $|S'| \leq 2(l - 2)$ . Thus  $S$  has at most  $1 + 2(l - 2) < 2(l - 1)$  edges.
2. Otherwise, partition the vertices adjacent to edges in  $S$  into two sets  $V_1$  and  $V_2$ , where  $V_1$  contains all vertices of degree one, and  $V_2$  contains all vertices of degree larger than one.




Figure 2: The bipartite graph  $G_n$  for  $f_n$ .

It is noted that since no length-3 paths exist, we have that  $|S| = |V_1|$ . If  $|V_2| \geq 2$ , then  $|V_1| \leq 2l - 2 = 2(l - 1)$ , since there are at most  $2l$  vertices adjacent to edges in  $S$ . In case  $|V_2| = 1$ , then  $S$  is a star, i.e. a single vertex  $u$  connected to a collection of vertices  $v_1, v_2, \dots, v_k$ . Then  $k \leq l$  and  $|S| = k \leq l \leq 2(l - 1)$ , for  $l \geq 2$ .

□

**Theorem 6.**  $f_n$  can not be represented as a sum of  $\lfloor n/2 \rfloor$  RO-formulas.

*Proof.* For the purpose of contradiction, suppose  $f_n$  can be represented as a sum of  $\lfloor n/2 \rfloor$  RO-formula-polynomials  $q_1, q_2, \dots, q_{\lfloor n/2 \rfloor}$ . Let  $G_n = (V_n, E_n)$  be the graph associated with  $f_n$ . For any  $q_i$ , let  $S_i \subseteq E_n$  be the set of edges representing the terms appearing in  $q_i$  of the form  $x_a x_b$ , where  $a \in [2n]$  is even, and  $b \in [2n]$  is odd. Note that since  $f$  has  $n^2$  many terms, some  $q_i$  should have  $|S_i| \geq 2n$ . Then by Claim 3,  $S_i$  contains a length-3 path. Therefore  $\alpha x_i x_j + \beta x_j x_k + \gamma x_k x_l$  appears in  $q_i$ , for distinct  $i, j, k$  and nonzero constants  $\alpha, \beta, \gamma \in \mathbb{F}$ . Due to Claim 2,  $q_i$  can not be computed by a RO-formula, which is a contradiction. □

## 4 Proof of Generator Lemma 1

Let  $p = |\text{Var}(f)|$ . The proof proceeds by induction on  $p$ . The bases  $p = 0$  and  $p = 1$  trivially hold.

Suppose  $p > 1$ . Hence  $m \geq 1$ . Consider arbitrary RO-ABP  $A$  computing  $f$ . Let  $s$  and  $t$  be the source and sink of  $A$ , respectively. Wlog. assume that only the  $p$  variables in  $\text{Var}(f)$  are present in  $A$ , and assume  $A$  satisfies the condition yielded by Lemma 2. Observe that for some variable  $x_i$  there are at most  $p/2$  variables in layers before the layer containing  $x_i$ , and at most  $p/2$  variables in layers after. (If  $p$  is odd it splits  $((p - 1)/2, (p - 1)/2)$  if  $p$  is even it splits  $(p/2 - 1, p/2)$ ).

Say  $x_i$  is on the arc  $(v_1, w_1)$  from layer  $L_j$  to  $L_{j+1}$ , where  $L_j = \{v_1, v_2, \dots, v_{m_1}\}$  and  $L_{j+1} = \{v_1, v_2, \dots, v_{m_2}\}$ , for certain  $j, m_1, m_2$ . We can write

$$f = \sum_{a=1}^{m_1} f_{s, v_a} f_{v_a, t}, \quad (2)$$

where for any nodes  $p$  and  $q$  in  $A$ ,  $f_{p,q}$  is the polynomial computed by subprogram of  $A_{p,q}$ . Consider  $f' = f(G_m^1, \dots, G_m^{i-1}, x_i, G_m^{i+1}, \dots, G_m^n)$ .

**Claim 1.** Write  $f' = x_i \cdot \frac{\partial f}{\partial x_i}(G_m^1, \dots, G_m^{i-1}, G_m^{i+1}, \dots, G_m^n) + f(G_m^1, \dots, G_m^{i-1}, 0, G_m^{i+1}, \dots, G_m^n)$ . Then  $\frac{\partial f}{\partial x_i}(G_m^1, \dots, G_m^{i-1}, G_m^{i+1}, \dots, G_m^n) \neq 0$ .

*Proof.* Since  $f$  depends on  $x_i$  and  $f$  is multilinear,  $\frac{\partial f}{\partial x_i} \neq 0$ . Let  $f'' = \frac{\partial f}{\partial x_i}$ . We will show that  $f''(G_m) \neq 0$ . Observe that in the r.h.s. of (2) only  $f_{v_1,t}$  depends on  $x_i$ . This implies that  $f'' = \frac{\partial f_{v_1,t}}{\partial x_i} \cdot f_{s,v_1}$ . Observe that  $|Var(f_{s,v_1})|$  and  $|Var(\frac{\partial f_{v_1,t}}{\partial x_i})|$  are both at most  $p/2$ . Since  $f'' \neq 0$ , both  $f_{s,v_1}$  and  $\frac{\partial f_{v_1,t}}{\partial x_i}$  are not identically zero. Certainly  $f_{s,v_1}$  can be computed by a RO-ABP. By Lemma 3, we know also  $\frac{\partial f_{v_1,t}}{\partial x_i}$  can be computed by a RO-ABP. As  $p/2 < p$ , the induction hypothesis applies. Since  $p/2 \leq 2^{m-1}$ , it yields that  $f_{s,v_1}(G_m) \neq 0$  and  $\frac{\partial f_{v_1,t}}{\partial x_i}(G_m) \neq 0$ . Therefore  $f''(G_m) \neq 0$ . This proves the claim.  $\square$

Recall the set  $A = \{a_1, \dots, a_n\}$  used for the construction of the SV-generator. By Observation 5.2 in [2],  $f(G_{m+1})|_{y_{m+1}=a_i} = f'|_{x_i=G_m^i+z_{m+1}}$ . Since  $z_{m+1}$  does not appear in  $G_m^j$  for any  $j$ , we get by Claim 1 that  $f(G_{m+1})|_{y_{m+1}=a_i} \neq 0$ . Hence  $f(G_{m+1}) \neq 0$ .  $\square$

## 5 X-Aligned RO-ABP-polynomials

The following lemma leads up to our central definition:

**Lemma 7.** . For all  $i \in [k]$ , Let  $f \in \mathbb{F}[X]$  be a RO-ABP-polynomial with  $|Var(f)| \geq 3$ . Then for any  $x_i \in Var(f)$ , there exist distinct  $x_j, x_k \in X \setminus \{x_i\}$  such that  $\frac{\partial^2 f}{\partial x_j \partial x_k} = g \cdot (\beta x_i - \alpha)$ , where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha, \beta \in \mathbb{F}$ .

*Proof.* Let  $A$  be a RO-ABP computing  $f$ . Wlog. assume all variables in  $X$  appear in  $A$ . By Lemma 2 assume wlog. that  $A$  has at most one variable per layer. Let  $x_{r_1}, x_{r_2}, \dots, x_{r_n}$  be the variables in  $X$  as they appear layer-by-layer, when going from the source to the sink of  $A$ . Consider an arbitrary  $x_i \in Var(f)$ . First, we handle the case that  $i = r_m$ , for some  $1 < m < n$ .

Let  $j = r_{m-1}$  and  $k = r_{m+1}$ . So  $x_j$  and  $x_k$  are the variables right before and right after  $x_i$  in  $A$ , respectively. Assume that  $x_j$  and  $x_k$  label the edges  $(u, v)$  and  $(m, n)$  respectively. Then  $\frac{\partial^2 f}{\partial x_j \partial x_k} = f_{s,u} f_{v,m} f_{n,t}$ , where  $f_{s,u} f_{v,m}$ , and  $f_{n,t}$  are computed by the subprograms  $A_{s,u}$ ,  $A_{v,m}$ , and  $A_{n,t}$ , respectively. Observe that  $f_{v,m}$  is of form  $\beta x_i - \alpha$ , for  $\alpha, \beta \in \mathbb{F}$ . Take  $g = f_{s,u} f_{v,m}$ , which is easily seen to be RO-ABP-computable by putting  $A_{s,u}$  and  $A_{v,m}$  in series, or by appealing to Lemmas 3 and 4.

The special case where  $i = r_1$  ( $i = r_n$ ), i.e.  $x_i$  is the first (last) variable in  $A$ , is handled similarly as above, by choosing  $x_k \in X \setminus \{x_i, x_j\}$  arbitrarily and appealing to Lemma 3.  $\square$

In the above lemma we have no guarantee the  $\alpha$  is nonzero, in case  $\beta \neq 0$ . We would like to consider polynomials which are in general position in this regard. We make the following definition:

**Definition 2.** Let  $S \subseteq X$ . Every RO-ABP-polynomial  $f \in \mathbb{F}[X]$  with  $|Var(f)| \leq 2$  is  $X$ -pre-aligned on  $S$ . A RO-ABP-polynomial  $f \in \mathbb{F}[X]$  with  $|Var(f)| > 2$  is  $X$ -pre-aligned on  $S$ , if the following condition is satisfied:

1. for every  $x_i \in S$ , there exist distinct  $x_j, x_k \in X \setminus \{x_i\}$  such that  $\frac{\partial^2 f}{\partial x_j \partial x_k} = g \cdot (\beta x_i - \alpha)$ , where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha, \beta \in F$  satisfy that  $\alpha = 0 \Rightarrow \beta = 0$ .

If  $f$  is  $X$ -pre-aligned on  $\text{Var}(f)$ , we simply say that  $f$  is  $X$ -pre-aligned.

For the  $X$ -pre-alignment property to hold recursively w.r.t. setting variables to zero, is a particularly desirable property of a RO-ABP-polynomial to have, as we will see. We make the following inductive definition:

**Definition 3.** Every RO-ABP-polynomial  $f \in \mathbb{F}[X]$  with  $|\text{Var}(f)| \leq 2$  is  $X$ -aligned. A RO-ABP-polynomial  $f \in \mathbb{F}[X]$  with  $|\text{Var}(f)| > 2$  is  $X$ -aligned, if the following conditions are satisfied:

1.  $f$  is  $X$ -pre-aligned, and
2. for every  $x_i \in \text{Var}(f)$ ,  $f|_{x_i=0}$  is  $X \setminus \{x_i\}$ -aligned.

Next we prove some of the needed properties of our notion, starting with the following easily verified statement:

**Proposition 4.** If  $f \in \mathbb{F}[X]$  is  $X$ -pre-aligned, then  $\forall \mu \in \mathbb{F}$ ,  $\mu \cdot f$  is  $X$ -pre-aligned. The same statement holds with aligned instead of pre-aligned.

The notion of  $X$ -pre-alignment is well-behaved w.r.t. taking partial derivatives. This will be crucial for obtaining the Hardness of Representation Theorem 8. We have the following lemma:

**Lemma 8.** For any RO-ABP-polynomial  $f \in \mathbb{F}[X]$  and any  $x_r \in X$ , the following hold:

1. If  $f$  is  $X$ -pre-aligned, then  $\frac{\partial f}{\partial x_r}$  is  $(X \setminus \{x_r\})$ -pre-aligned.
2. If  $f$  is  $X$ -aligned, then  $\frac{\partial f}{\partial x_r}$  is  $(X \setminus \{x_r\})$ -aligned.

*Proof.* We first show that Item 1 holds. Let  $f' = \frac{\partial f}{\partial x_r}$  and  $X' = X \setminus \{x_r\}$ . By Lemma 3, we know that  $f'$  is a RO-ABP-polynomial. Assume that  $|\text{Var}(f')| \geq 3$ , since otherwise the statement holds trivially. Consider arbitrary  $x_i \in \text{Var}(f')$ . Then  $x_i \in \text{Var}(f)$ , so there exist distinct  $x_j$  and  $x_k$  in  $X \setminus \{x_i\}$ , such that  $\frac{\partial^2 f}{\partial x_j \partial x_k} = g \cdot (\beta x_i - \alpha)$ , where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha = 0 \Rightarrow \beta = 0$ . Consider the following two cases:

**Case I:**  $r \notin \{j, k\}$ .

Hence  $x_j, x_k \in X' \setminus \{x_i\}$ . We have that  $\frac{\partial^2 f'}{\partial x_j \partial x_k} = \frac{\partial^3 f}{\partial x_j \partial x_k \partial x_r} = \frac{\partial g}{\partial x_r} \cdot (\beta x_i - \alpha)$ . By Lemma 3,  $\frac{\partial g}{\partial x_r}$  is a RO-ABP-polynomial, and it clearly does not depend on  $x_i$ , so we conclude that  $f'$  is  $X'$ -pre-aligned on  $\{x_i\}$ .

**Case II:**  $r \in \{j, k\}$ .

Wlog. assume  $r = j$ . Then  $x_k \in X' \setminus \{x_i\}$ . Since  $|\text{Var}(f')| \geq 3$ , there must be at least one more variable  $x_l$  in  $\text{Var}(f')$  distinct from each of  $x_k$  and  $x_i$ . Then  $x_l \in X' \setminus \{x_i\}$ . We have that  $\frac{\partial f'}{\partial x_k} = g \cdot (\beta x_i - \alpha)$ . Hence  $\frac{\partial^2 f'}{\partial x_k \partial x_l} = \frac{\partial g}{\partial x_l} \cdot (\beta x_i - \alpha)$ . We again conclude  $f'$  is  $X'$ -pre-aligned on  $\{x_i\}$ .

Since in the above,  $x_i$  was taken arbitrarily from  $\text{Var}(f')$ , we conclude  $f'$  is  $X'$ -pre-aligned.

Item 2 is proved by induction on  $|X|$ . The base case is when  $|X| \leq 3$ . Then  $|\text{Var}(f')| \leq 2$ , and hence  $f'$  is  $X'$ -aligned. Now suppose  $|X| > 3$ . Assume  $|\text{Var}(f')| > 2$ , since otherwise it is trivial. By Item 1, we know  $f'$  is  $X'$ -pre-aligned. Consider an arbitrary  $x_i \in \text{Var}(f')$ . Then  $x_i \in \text{Var}(f)$ .

We have that  $f'_{|x_i=0} = \left(\frac{\partial f}{\partial x_r}\right)_{x_i=0} = \frac{\partial f_{|x_i=0}}{\partial x_r}$ . Since  $f_{|x_i=0}$  is  $(X \setminus \{x_i\})$ -aligned, we can apply the induction hypothesis to conclude that  $\frac{\partial f_{|x_i=0}}{\partial x_r}$  is  $(X \setminus \{x_i\}) \setminus \{x_r\} = (X' \setminus \{x_i\})$ -aligned.  $\square$

## 5.1 A Workable Sufficient Condition

Next we establish a sufficient condition, so for a given RO-ABP-polynomial  $f$  we can make  $f(x_1 + v_1, x_2 + v_2, \dots, x_n + v_n)$   $X$ -aligned, by means of computing some shift  $v \in \mathbb{F}^n$ . For this, let us call a polynomial  $f \in \mathbb{F}[X]$  *decent*, if for all  $x_a, x_b \in \text{Var}(f)$  with  $\frac{\partial^2 f}{\partial x_a \partial x_b} \neq 0$ , it holds that the monomial  $x_a x_b$  appears in  $f$  with a nonzero constant coefficient.

**Lemma 9.** *A RO-ABP-polynomial  $f \in \mathbb{F}[X]$  is  $X$ -aligned, if  $|\text{Var}(f)| \leq 2$ , or else for any  $I \subseteq \text{Var}(f)$  with  $|I| \leq |\text{Var}(f)| - 3$ ,  $f_{|x_I=0}$  is decent.*

*Proof.* We use induction on  $|\text{Var}(f)|$ . For the base case  $|\text{Var}(f)| \leq 2$  it is trivial. Now assume  $|\text{Var}(f)| > 2$ . Take  $I = \emptyset$ . Then we get that for any  $x_a, x_b \in \text{Var}(f)$ , if  $\frac{\partial^2 f}{\partial x_a \partial x_b} \neq 0$  then the monomial  $x_a x_b$  appears in  $f$  with a nonzero constant coefficient.

Let us first establish that  $f$  is  $X$ -pre-aligned. Consider an arbitrary  $x_i \in \text{Var}(f)$ . By Lemma 7, there exist distinct  $x_j, x_k \in X \setminus \{x_i\}$  such that

$$\frac{\partial^2 f}{\partial x_j \partial x_k} = g \cdot (\beta x_i - \alpha), \quad (3)$$

where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha, \beta \in F$ .

If  $\beta = 0$ , then  $f$  is  $X$ -pre-aligned on  $\{x_i\}$ , so suppose  $\beta \neq 0$ . If (3) is identically zero, then we know  $g \equiv 0$ , so  $\frac{\partial^2 f}{\partial x_j \partial x_k} = g \cdot (\beta x_i - \alpha')$ , for any arbitrary  $\alpha' \neq 0$ . If (3) is not identically zero, then we know  $x_j x_k$  is in  $f$ , which implies that  $\alpha \neq 0$ . We conclude that  $f$  is  $X$ -pre-aligned on  $\{x_i\}$ .

In the above, we find that  $f$  is  $X$ -pre-aligned on  $\{x_i\}$  in any of the considered cases. Since  $x_i$  was arbitrarily taken from  $\text{Var}(f)$ , we conclude that  $f$  is  $X$ -pre-aligned.

Next, we show Condition 2 of Definition 3 holds. Consider  $f' := f_{|x_i=0}$ , for an arbitrary  $x_i \in \text{Var}(f)$ . We want to establish that the sufficient condition of Lemma 9 holds for  $f' \in \mathbb{F}[X \setminus \{x_i\}]$ , since then we can by apply the induction hypothesis and conclude that  $f'$  is  $(X \setminus \{x_i\})$ -aligned.

If  $|\text{Var}(f')| \leq 2$  the sufficient condition of the Lemma 9 clearly holds for  $f'$ . Otherwise, consider  $I' \subseteq \text{Var}(f')$  of size at most  $|\text{Var}(f')| - 3$ . Let  $I = I' \cup \{x_i\}$ . Then  $|I| \leq |\text{Var}(f)| - 3$ . Now consider  $x_a, x_b \in \text{Var}(f'_{|x_{I'}=0}) = \text{Var}(f_{|x_I=0})$ . Suppose  $\frac{\partial^2 f'_{|x_{I'}=0}}{\partial x_a \partial x_b} \neq 0$ . Since the latter equals  $\frac{\partial^2 f_{|x_I=0}}{\partial x_a \partial x_b} \neq 0$ , we know that  $x_a x_b$  appears with a nonzero constant coefficient in  $f_{|x_I=0}$ . This implies  $x_a x_b$  appears with a nonzero constant coefficient in  $f_{|x_{I'}=0}$ . Hence  $f'_{|x_{I'}=0}$  is decent.

We conclude the sufficient condition of the Lemma 9 holds for  $f' \in \mathbb{F}[X \setminus \{x_i\}]$ . Hence by the induction hypothesis we conclude that  $f'$  is  $(X \setminus \{x_i\})$ -aligned.  $\square$

**Lemma 10.** *Any decent RO-ABP-polynomial  $f \in \mathbb{F}[X]$  is  $X$ -aligned.*

*Proof.* We show that the condition of Lemma 9 is satisfied. If  $|\text{Var}(f)| \leq 2$  this is clear. Otherwise, consider arbitrary  $I \subseteq \text{Var}(f)$  with  $|I| \leq |\text{Var}(f)| - 3$ . Let  $x_a, x_b \in \text{Var}(f_{|x_I=0})$ , be such that  $\frac{\partial^2 f_{|x_I=0}}{\partial x_a \partial x_b} \neq 0$ . We have that  $x_a, x_b \in \text{Var}(f)$ , and it must be that  $\frac{\partial^2 f}{\partial x_a \partial x_b} \neq 0$ , since  $\frac{\partial^2 f_{|x_I=0}}{\partial x_a \partial x_b} = \left(\frac{\partial^2 f}{\partial x_a \partial x_b}\right)_{|x_I=0}$ . Hence  $x_a x_b$  is in  $f$ . This implies that  $x_a x_b$  is in  $f_{|x_I=0}$ .  $\square$

## 5.2 Nearly Unique Nonalignment

In addition to the above, we crucially need the following “Nearly Unique Nonalignment Lemma”.

**Lemma 11.** *Let  $f \in \mathbb{F}[X]$  be an  $X$ -pre-aligned RO-ABP-polynomial for which  $\frac{\partial^2 f}{\partial x_p \partial x_q} \neq 0$ , for any distinct  $x_p, x_q \in X$ . Then there are at most two  $\gamma \in \mathbb{F}$  such that  $f|_{x_n=\gamma}$  is not  $(X \setminus \{x_n\})$ -pre-aligned.*

Before giving the proof, we need a lemma.

**Lemma 12.** *Let  $f \in \mathbb{F}[X]$  be a RO-ABP-polynomial with  $|\text{Var}(f)| \geq 3$  that is  $X$ -pre-aligned on  $S$ , for some  $S \subseteq \text{Var}(f)$ . Assume that for any distinct  $x_p, x_q \in X$ ,  $\frac{\partial^2 f}{\partial x_p \partial x_q} \neq 0$ . In any RO-ABP  $A$  computing  $f$ , for any  $x_i \in S$ ,*

1. *if there exists a non-constant layer with variable  $x_a$  right before the  $x_i$ -layer, and there exists a non-constant layer with variable  $x_b$  right after the  $x_i$ -layer, then*

$$\frac{\partial^2 f}{\partial x_a \partial x_b} = g \cdot (\beta x_i - \alpha),$$

*where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha, \beta \in F$  satisfy that  $\alpha = 0 \Rightarrow \beta = 0$ . Furthermore,  $-\alpha$  equals the sum of weights of all paths from  $\text{end}(x_a)$  to  $\text{begin}(x_b)$  that do not go over  $x_i$ .*

*Proof.* Consider  $x_i \in S$ . Since  $f$  is  $X$ -pre-aligned on  $S$ , we know there exist distinct  $x_j, x_k \in X \setminus \{x_i\}$  with  $\frac{\partial^2 f}{\partial x_j \partial x_k} = h \cdot (\beta' x_i - \alpha')$ , where  $h$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha', \beta' \in F$  satisfy that  $\alpha' = 0 \Rightarrow \beta' = 0$ . Since  $\frac{\partial^2 f}{\partial x_j \partial x_k} \neq 0$ , it must be that  $\alpha' \neq 0$ .

**Case I:** In  $A$ , the  $x_i$ -layer lies in between the  $x_j$ -layer and  $x_k$  layer.

Wlog assume the  $x_i$  layer lies before the  $x_k$ -layer and after the  $x_j$ -layer (according to the order of the DAG underlying  $A$ ). Write  $\frac{\partial^2 f}{\partial x_j \partial x_k} = p_1 p_2 \cdot (q_1 q_2 x_i + q_3)$ , where

- $p_1$  is the sum of weights over all paths in  $A$  from  $\text{source}(A)$  to  $\text{begin}(x_j)$ , and  $p_2$  is the sum of weights over all paths in  $A$  from  $\text{end}(x_k)$  to  $\text{sink}(A)$ .
- $q_3$  is the sum of weights over all paths from  $\text{end}(x_j)$  to  $\text{begin}(x_k)$  that bypass the  $x_i$ -edge,  $q_1$  is the sum of weights over all paths from  $\text{end}(x_j)$  to  $\text{begin}(x_i)$ , and  $q_2$  is the sum of weights over all paths from  $\text{end}(x_i)$  to  $\text{begin}(x_k)$ .

Now we have that  $p_1 p_2 \cdot (q_1 q_2 x_i + q_3) = h \cdot (\beta' x_i - \alpha')$ . Since both  $p_1 p_2$  and  $h$  do not depend on  $x_i$ , it must be that  $(\beta' x_i - \alpha') \mid (q_1 q_2 x_i + q_3)$ . Note that  $\beta'$  cannot equal 0, since then one of  $q_1, q_2$  would be zero. The latter implies that  $\frac{\partial^2 f}{\partial x_i \partial x_j} \equiv 0$  or  $\frac{\partial^2 f}{\partial x_i \partial x_k} \equiv 0$ , which is a contradiction. Since  $\beta' \neq 0$ , we can conclude that  $q_3 = \mu q_1 q_2$  for some  $\mu \in \mathbb{F}$ ,  $\mu \neq 0$ . Now we need the following claim:

**Claim 2.** *Given an RO-ABP  $A$  computing  $f(x_1, \dots, x_n)$ , if for any distinct  $x_p, x_q \in X$ ,  $\frac{\partial^2 f}{\partial x_p \partial x_q} \neq 0$ , then  $\prod_{i \in [n]} x_i$  appears in  $f$ . Furthermore, for two variables  $x_i$  and  $x_j$ , if  $x_i$  is before  $x_j$  in  $A$ , if we let  $S$  be the set of variables in between  $x_i$  and  $x_j$ , then  $\prod_{x_m \in S} x_m$  is a term in the polynomial  $\hat{A}(\text{end}(x_i), \text{begin}(x_j))$ .*

*Proof.* Suppose the variable layers in  $A$  are arranged according to the permutation  $\phi : [n] \rightarrow [n]$ , that is,  $x_{\phi(i)}$  labels the  $i$ th variable layer. Then we that

1.  $\hat{A}(s, \text{begin}(x_{\phi(1)})) \neq 0$  (Since otherwise  $\frac{\partial^2 f}{\partial x_{\phi(1)} \partial x_{\phi(2)}} \equiv 0$ ),
2. Similarly  $\hat{A}(\text{end}(x_{\phi(n)}), t) \neq 0$ , and
3. For  $i \in [n-1]$ ,  $\hat{A}(\text{begin}(x_{\phi(i)}), \text{end}(x_{\phi(i+1)})) \neq 0$  (Since otherwise  $\frac{\partial^2 f}{\partial x_{\phi(i)} \partial x_{\phi(i+1)}} \equiv 0$ ).

The coefficient of  $\prod_{i \in [n]} x_i$  is just

$$\hat{A}(s, \text{begin}(x_{\phi(1)})) \cdot \hat{A}(\text{end}(x_{\phi(n)}), t) \prod_{i \in [n-1]} \hat{A}(\text{begin}(x_{\phi(i)}), \text{end}(x_{\phi(i+1)})),$$

and hence  $\prod_{i \in [n]} x_i$  appears in  $f$ . A similar argument yields the statement for  $\hat{A}(\text{end}(x_i), \text{begin}(x_j))$ .  $\square$

As in the proof of Lemma 7, write  $\frac{\partial^2 f}{\partial x_a \partial x_b} = g \cdot (\beta x_i - \alpha)$ , where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $-\alpha$  equals the sum of weights over all paths from  $\text{end}(x_a)$  to  $\text{begin}(x_b)$  not going over  $x_i$ . We have three cases:

1. Neither  $x_j$  nor  $x_k$  is the most adjacent variable to  $x_i$  in  $A$ . By above claim,  $x_a$  appears in a monomial of  $q_1$ , and  $x_b$  appears in a monomial  $q_2$ . Hence, there is a monomial in  $q_1 q_2$  with  $x_a x_b$ . As  $q_3 = \mu q_1 q_2$ , for  $\mu \neq 0$ , the same can be said for  $q_3$ . But this implies  $\alpha \neq 0$ , as the coefficient of  $x_a x_b$  is  $-\alpha \cdot \hat{A}(\text{end}(x_j), \text{begin}(x_a)) \hat{A}(\text{end}(x_b), \text{begin}(x_k))$ .
2.  $x_j$  is not the most adjacent variable to  $x_i$  in  $A$ , but  $x_k = x_b$ . Then similarly  $q_1 q_2$  has a monomial with  $x_a$  in it, and therefore the same holds for  $q_3$ . Therefore  $\alpha \neq 0$ , as the coefficient of  $x_a$  in  $q_3$  is  $-\alpha \cdot \hat{A}(\text{end}(x_j), \text{begin}(x_a))$ .
3.  $x_j = x_a$ , but  $x_k$  is not the most adjacent variable to  $x_i$  in  $A$ . This is argued similarly as the second item.

This concludes the argument for this case.

**Case II:** In  $A$ , the  $x_i$ -layer lies before the  $x_j$ -layer and  $x_k$ -layer.

Wlog. assume that the  $x_j$  layer lies before the  $x_k$  layer. Similarly as in Case I, we write  $\frac{\partial^2 f}{\partial x_j \partial x_k} = p_1 p_2 \cdot (q_1 q_2 x_i + q_3)$ , but where now we have that

- $p_1 = \hat{A}_{\text{end}(x_j), \text{begin}(x_k)}$ , and  $p_2 = \hat{A}_{\text{end}(x_k), \text{sink}(A)}$ ,
- $q_1 = \hat{A}_{\text{source}(A), \text{begin}(x_i)}$ ,
- $q_2 = \hat{A}_{\text{end}(x_i), \text{begin}(x_j)}$ ,
- $q_3 = A[x_i = 0]_{\text{source}(A), \text{begin}(x_j)}$ .

Then  $p_1 p_2 \cdot (q_1 q_2 x_i + q_3) = h \cdot (\beta' x_i - \alpha')$ . Since both  $p_1 p_2$  and  $h$  do not depend on  $x_i$ , it must be that  $(\beta' x_i - \alpha') \mid (q_1 q_2 x_i + q_3)$ . Similarly as before, we get  $q_3 = \mu q_1 q_2$  for some  $\mu \in \mathbb{F}$ ,  $\mu \neq 0$ .

The rest of the proof is similar to Case I. One argues that 1) when  $x_j \neq x_b$ ,  $q_1 q_2$  contains a monomial with  $x_a x_b$ . To make  $x_a x_b$  appear in a monomial  $q_3$  we need  $\alpha \neq 0$ , and 2) when  $x_j = x_b$ ,  $q_1 q_2$  contains a monomial with  $x_a$ , and to make  $x_a$  appear in a monomial of  $q_3$ , we need  $\alpha \neq 0$ .

**Case III:** In  $A$ , the  $x_i$ -layer lies after the  $x_j$ -layer and  $x_k$ -layer.

This case is symmetrical to Case II.  $\square$

We also need the following proposition:

**Proposition 5.** *Let  $f \in \mathbb{F}[X]$  be a RO-ABP-polynomial with  $|Var(f)| \geq 3$ , and let  $S \subseteq Var(f)$ . Then  $f$  is  $X$ -pre-aligned on  $S$  if and only if  $f' := (x_{n+1} + 1)f$  is  $X \cup \{x_{n+1}\}$ -pre-aligned on  $S$ .*

*Proof.* Let  $X' = X \cup \{x_{n+1}\}$ . It is easy to see that assuming  $f$  is  $X$ -pre-aligned on  $S$ , we have that  $f$  is  $X'$ -pre-aligned on  $S$ .

Conversely, assume  $f'$  is  $X'$ -pre-aligned on  $S$ . Let  $x_i \in S$ . Then there exist  $x_j, x_k \in X' \setminus \{x_i\}$ , such that  $\frac{\partial^2 f'}{\partial x_j \partial x_k} = g(\beta x_i + \alpha)$ , where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha = 0$  implies  $\beta = 0$ . If  $x_{n+1} \notin \{x_j, x_k\}$ , then  $\frac{\partial^2 f'}{\partial x_j \partial x_k} = \frac{\partial^2 f}{\partial x_j \partial x_k}(x_{n+1} + 1)$ . Setting  $x_{n+1} = 0$ , we have that  $\frac{\partial^2 f}{\partial x_j \partial x_k} = (g|_{x_{n+1}=0})(\beta x_i + \alpha)$ . So we get the required  $X$ -pre-alignment of  $f$  on  $\{x_i\}$ . Otherwise, say wlog.  $x_j = x_{n+1}$ . We have that  $\frac{\partial f}{\partial x_k} = \frac{\partial^2 f'}{\partial x_{n+1} \partial x_k} = g(\beta x_i + \alpha)$ . One easily obtains the required  $X$ -pre-alignment of  $f$  on  $\{x_i\}$ , by taking one more  $\partial x_l$ , for some variable  $x_l \in X \setminus \{x_i, x_k\}$ , and then using Lemma 3.  $\square$

We are now ready to give the proof of Lemma 11.

### 5.3 Proof

We prove the lemma by induction on  $|X|$ . For the base case we take  $|X| \leq 3$ , in which case the statement clearly holds. Now suppose  $|X| > 3$ . Let  $f' = f|_{x_n=\gamma}$ , for some  $\gamma$ . Let  $X' = X \setminus \{x_n\}$ . Suppose  $f'$  is not  $X'$ -pre-aligned. Hence  $|Var(f')| \geq 3$ . We want to show this can happen for at most one  $\gamma$ .

Consider an arbitrary RO-ABP  $A$  computing  $f$ . Let  $f_e = f(x_{n+1}+1)(x_{n+2}+1)(x_{n+3}+1)(x_{n+4}+1)$ . Let  $X_e := X \cup \{x_{n+1}, x_{n+2}, x_{n+3}, x_{n+4}\}$ . By Proposition 5,  $f_e$  is  $X_e$ -pre-aligned on  $Var(f)$ . Let  $f'_e := (f_e)|_{x_n=\gamma}$  and  $X'_e := X' \cup \{x_{n+1}, x_{n+2}, x_{n+3}, x_{n+4}\}$ . Note that  $f'_e = f'(x_{n+1}+1)(x_{n+2}+1)(x_{n+3}+1)(x_{n+4}+1)$ . So also by Proposition 5,  $f'_e$  is not  $X'_e$ -pre-aligned on  $Var(f')$  if and only if  $f'$  is not  $X'$ -pre-aligned on  $Var(f')$ . We will show the former happens for at most one  $\gamma$ . So let us assume that  $f'_e$  is not  $X'_e$ -pre-aligned on  $Var(f')$ . We can easily obtain a RO-ABP  $A_e$  from  $A$ , which computes  $f_e$ . In this, we make sure  $x_{n+1}$  and  $x_{n+2}$  are the first and second variable in  $A_e$ , and  $x_{n+3}$  and  $x_{n+4}$  are the fore-last and last variable in  $A_e$ . For each  $x_i \in Var(f')$ , let  $x_{j_i}$  be the variable right after  $x_i$  in  $A_e$ , and let  $x_{k_i}$  be the variable before  $x_i$  in  $A_e$ . Note that we have made sure these always exist in  $A_e$ . Since  $f_e$  is  $X_e$ -pre-aligned on  $Var(f)$ , by Lemma 12,  $\frac{\partial^2 f_e}{\partial x_{j_i} \partial x_{k_i}} = g \cdot (\beta_i x_i - \alpha_i)$ , where  $g$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha_i = 0 \Rightarrow \beta_i = 0$ . Furthermore, we have that  $\alpha_i$  is the sum of weights of all paths from  $end(x_{k_i})$  to  $begin(x_n)$ , which do not go over  $x_i$  in  $A_e$ . Consider the following two cases:

**Case I:**  $n \notin \{j_i, k_i\}$ , for any  $x_i \in Var(f')$ .

Then for any  $i$ ,  $\frac{\partial^2 f'_e}{\partial x_{j_i} \partial x_{k_i}} = (g_i)|_{x_n=\gamma} \cdot (\beta_i x_i - \alpha_i)$ , which contradicts the assumption that  $f'_e$  is not  $X'_e$ -pre-aligned on  $Var(f')$ .

**Case II:**  $n \in \{j_i, k_i\}$ , for some  $x_i \in Var(f')$ .

By symmetry we can assume wlog. that  $j_i = n$  (the case  $k_i = n$  is handled similarly). Since  $\frac{\partial^2 f}{\partial x_{j_i} \partial x_{k_i}} \not\equiv 0$ , and  $\alpha_i = 0$  implies  $\beta_i = 0$ , We have that  $\alpha_i \neq 0$ .

We know that in  $A_e$  there still exists a variables layer, say with variables  $x_l$ , right after the  $x_{j_i}$ -layer. Let  $b_i = \text{begin}(x_i)$ ,  $e_i = \text{end}(x_i)$ ,  $b_n = \text{begin}(x_n)$ , and  $e_n = \text{end}(x_n)$ . Let  $s = \text{end}(x_{k_i})$  and  $t = \text{begin}(x_l)$ . Then write:

$$\frac{\partial^2 f_e}{\partial x_l \partial x_{k_i}} = p_1 p_2 (c_{s,b_i} c_{e_i,b_n} c_{e_n,t} x_i x_n + c_{s,b_i} c_{e_i,t} x_i + c_{s,b_n} c_{e_n,t} x_n + c_{s,t}),$$

where in the above each constant  $c_{v,w}$  is the sum of weights over all paths from  $v$  to  $w$  going over constant labeled edges only. Note that  $c_{s,b_n} = \alpha_i \neq 0$ . Furthermore,  $p_1$  is the sum of weights of all paths from  $\text{source}(A_e)$  to  $\text{begin}(x_{k_i})$ , and  $p_2$  is the sum of weights over all paths from  $\text{end}(x_l)$  to  $\text{sink}(A_e)$ . Then

$$\frac{\partial^2 f'_e}{\partial x_l \partial x_{k_i}} = p_1 p_2 ((c_{s,b_i} c_{e_i,b_n} c_{e_n,t} \gamma + c_{s,b_i} c_{e_i,t}) x_i + c_{s,b_n} c_{e_n,t} \gamma + c_{s,t}),$$

We have that  $f'_e$  can only not be  $X'_e$ -pre-aligned on  $\{x_i\}$  if  $c_{s,b_n} c_{e_n,t} \gamma + c_{s,t} = 0$ . This can happen for more than one  $\gamma$  only if  $c_{s,b_n} c_{e_n,t} = 0$ . Since  $c_{s,b_n} \neq 0$ , this happens only if  $c_{e_n,t} = 0$ , but the latter implies that  $\frac{\partial^2 f_e}{\partial x_l \partial x_n} \equiv 0$ , which in turn implies that  $\frac{\partial^2 f}{\partial x_l \partial x_n} \equiv 0$ , which is a contradiction.

Finally, putting together from what we observed from the above two cases, note that, Case II can apply at most twice for a variable  $x_i \in \text{Var}(f')$ . Namely, possibly once for the variable right before  $x_n$ , and possibly once for the variable after  $x_n$ . We conclude the lemma holds.  $\square$

**Corollary 1.** *Suppose  $|\mathbb{F}| > 3$ . Let  $h, g \in \mathbb{F}[X]$  be RO-ABP-polynomials such that  $h = g \cdot (\beta x_n - \alpha)$ , for  $\beta \in \mathbb{F} \setminus \{0\}$ . If  $h$  is  $X$ -pre-aligned, then  $g$  is  $(X \setminus \{x_n\})$ -pre-aligned.*

*Proof.* If we set  $x_n$  to any value  $\gamma \neq \alpha/\beta$ , we get that  $h|_{x_n=\gamma}$  is a nonzero constant multiple of  $g$ . By Lemma 11, there are at most two  $\gamma$  such that  $h|_{x_n=\gamma}$  is not  $(X \setminus \{x_n\})$ -pre-aligned. Now use Proposition 4 to conclude that  $g$  is  $(X \setminus \{x_n\})$ -pre-aligned.  $\square$

## 6 Simultaneous Alignment of RO-ABP-polynomials

**Definition 4.** *A simultaneous  $X$ -alignment for a set of RO-ABP-polynomials  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$  is a vector  $v \in \mathbb{F}^n$  such that  $f_i(x_1 + v_1, x_2 + v_2, \dots, x_n + v_n)$  is  $X$ -aligned for every  $i \in [k]$ .*

We present an algorithm for finding a simultaneous  $X$ -alignment for a set of RO-ABP-polynomials. We assume that we have a polynomial identity testing algorithm  $\text{PIT}_{\text{RO-ABP}}$  for testing a single RO-ABP. We prove a corollary of Lemma 10 first.

**Corollary 2.** *Let  $\{f_i\}_{i \in [k]}$  be a set of RO-ABP-polynomials in  $\mathbb{F}[X]$ . Then  $v \in \mathbb{F}^n$  is a simultaneous  $X$ -alignment for  $\{f_i\}_{i \in [k]}$ , if it is a simultaneous nonzero for  $\{\frac{\partial^2 f_i}{\partial x_a \partial x_b} \mid \frac{\partial^2 f_i}{\partial x_a \partial x_b} \neq 0\}_{i \in [k], a, b \in [n]}$ .*

*Proof.* Consider  $\{f'_i = f_i(x_1 + v_1, x_2 + v_2, \dots, x_n + v_n)\}_{i \in [k]}$ . Due to Lemma 10, we only need to show that for every  $i$ , for every  $x_a, x_b \in \text{Var}(f_i)$ , if  $\frac{\partial^2 f'_i}{\partial x_a \partial x_b} \neq 0$  then the monomial  $x_a x_b$  appears in  $f'_i$  with a nonzero constant coefficient. Observe that the monomial  $x_a x_b$  appears in  $f'_i$  with a nonzero constant coefficient  $\iff \frac{\partial^2 f'_i}{\partial x_a \partial x_b}(\bar{0}) \neq 0$ . The latter holds, as  $\frac{\partial^2 f'_i}{\partial x_a \partial x_b}(\bar{0}) = \frac{\partial^2 f_i}{\partial x_a \partial x_b}(v) \neq 0$ .  $\square$

Now the argument is similar as for Lemma 4.3 in [2], but with first order partial derivatives replaced by second order ones. This yields the following theorem:

**Theorem 7.** *Let  $\mathbb{F}$  be a field with  $|\mathbb{F}| > kn^2$ . There exists an algorithm for finding a simultaneous  $X$ -alignment for a set of RO-ABP polynomials  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$ . The algorithm makes oracle calls to the procedure  $\text{PIT}_{\text{RO-ABP}}$ . The  $f_i$ s are only accessed through this subroutine. The running-time of the algorithm is  $O(k^2 n^5 \cdot t)$ , where  $t$  is an upper bound on the time needed for any subroutine call to  $\text{PIT}_{\text{RO-ABP}}$ .*

*Proof.* We assume that we have a polynomial identity testing algorithm  $\text{PIT}_{\text{RO-ABP}}$  for testing a single RO-ABP, such that  $\text{PIT}_{\text{RO-ABP}}$  outputs *True* if  $f \equiv 0$  and *False* otherwise. We have the following algorithm:

---

**Algorithm 1** Alignment Finding.

---

**Input:** A set of RO-ABP-polynomials  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$ .

**Output:** A simultaneous alignment  $v$  for  $\{f_i\}_{i \in [k]}$ .

**Oracle:** PIT algorithm  $\text{PIT}_{\text{RO-ABP}}$ .

```

1:  $L = \emptyset$ 
2: for all  $f_i$  and  $(x_a, x_b)$ ,  $a, b \in [n]$ ,  $a \neq b$  do
3:   If  $\text{PIT}_{\text{RO-ABP}}(\frac{\partial^2 f_i}{\partial x_a \partial x_b}) = \text{False}$ , add it to  $L$ 
4: end for
5: for all  $j \in [n]$  do
6:   Find  $c$  such that for every  $g \in L$ ,  $\text{PIT}_{\text{RO-ABP}}(g|_{x_j=c}) = \text{False}$ 
7:    $v_j \leftarrow c$ 
8:   For every  $g \in L$ ,  $g \leftarrow g|_{x_j=c}$ 
9: end for
10: return  $v$ 

```

---

We first make two remarks, which pertain to applying Algorithm 1 in the setting where we only have black-box access to each  $f_i$ . Consider the set  $L$  the algorithm constructs with the execution of the first **for**-loop. Since we only have black-box access to  $f_i$ , the given pseudocode is intended to mean  $L$  is constructed symbolically. Having black-box access to  $f_i$  is enough to have black-box access to any element of  $L$ . Namely, by Lemma 3,  $f' := \frac{\partial^2 f_i}{\partial x_a \partial x_b}$  is a RO-ABP. Note that black-box access to  $f_i$  is sufficient for being able to compute  $f'(a)$  for any  $a \in \mathbb{F}^n$ . This is all the black-box RO-ABP algorithm needs to decide whether  $f' \equiv 0$ .

Similarly, on line 8 the substitution is not actually carried out, but done symbolically. So it is just remembered that  $x_j$  is set to  $c$ . For example, suppose that up to some point in the execution the algorithm it has set  $x_i = c_i$ , for  $i \in [m]$ . Then on line 6, for evaluating  $\text{PIT}_{\text{RO-ABP}}(g|_{x_j=c})$ , the black-box algorithm is granted access to a RO-ABP in  $n - m$  variables  $g(c_1, c_2, \dots, c_m, x_{m+1}, \dots, x_n)$ . The queries it makes can be answered with only black-box access to  $g$ .

Now, by Corollary 2 it suffices to find a common nonzero of the set  $L$ . First however, we need to explain how to find  $c$  such that  $g|_{x_j=c} \neq 0$ . Let  $V \subset \mathbb{F}$  with  $|V| = kn^2 + 1$  be given. We claim  $V$  always includes a good value. This is because we have at most  $kn^2$  multilinear polynomials in  $L$ , and for a specific one there is at most one bad value, due to Lemma 6. The algorithm can simply try all elements in  $V$  to get the required  $c$ . The correctness of the algorithm is now evident, from the observation that it simply maintains the invariant that all  $g \in L$  are not identically zero.

The running time of the algorithm is as follows: for line 2 we need  $O(kn^2)$  calls to  $\text{PIT}_{\text{RO-ABP}}$ . For line 7 we need  $O(n \cdot (kn^2 + 1) \cdot (kn^2)) = O(k^2n^5)$  calls to  $\text{PIT}_{\text{RO-ABP}}$ . Thus the total running time of the algorithm is  $O(k^2n^5 \cdot t)$ , where  $t$  is an upper bound on the time needed for any subroutine call to  $\text{PIT}_{\text{RO-ABP}}$ .  $\square$

By Lemma 1 and using Lemma 5,  $\text{PIT}_{\text{RO-ABP}}$  can be implemented in the black-box setting to run in time  $n^{O(\log n)}$ , where  $n$  is the number of variables of the input RO-ABP-polynomial. In the non-black-box setting, as is show in Appendix C,  $\text{PIT}_{\text{RO-ABP}}$  can be implemented to run in time  $O(n^2s)$ , when given an RO-ABP over  $n$  variables of size  $s$ . This yields the following two corollaries:

**Corollary 3.** *Provided  $|\mathbb{F}| > kn^2$ , there exists a non-black-box algorithm for finding a simultaneous  $X$ -alignment for a set  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$ , where  $f_i$  is computed by a RO-ABP  $A_i$ , for  $i \in [k]$ . The algorithm receives  $\{A_i\}_{i \in [k]}$  on the input, and it runs in time  $O(k^2n^7s)$ , where  $s$  is an upper bound on the size of any  $A_i$ .*

**Corollary 4.** *Provided  $|\mathbb{F}| > kn^2$ , there exists a black-box algorithm for finding a simultaneous  $X$ -alignment for a set of RO-ABP-polynomials  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$ . The algorithm queries individual  $f_i$ s, and runs in time  $k^2n^{O(\log n)}$ .*

## 6.1 Simultaneous Alignment Hitting Set

Here we present a black-box algorithm to find a candidate set  $\mathcal{A}_k$  of size  $(kn)^{O(\log n)}$ , which is guaranteed to contain a simultaneous  $X$ -alignment for any set of  $k$  RO-ABP-polynomials  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$ .

**Lemma 13.** *Let  $\mathbb{F}$  be a field with  $|\mathbb{F}| > kn^4$ , and let  $V \subseteq \mathbb{F}$  with  $|V| = kn^4 + 1$  be given. Let  $\{f_i\}_{i \in [k]}$  be a set of RO-ABP-polynomials in  $\mathbb{F}[X]$ . Let  $G_m : \mathbb{F}^{2m} \rightarrow \mathbb{F}^n$  be the  $m$ th-order SV-generator with  $m = \lceil \log n \rceil + 1$ . Then  $\mathcal{A}_k := G_m(V^{2m})$  contains a simultaneous  $X$ -alignment for  $\{f_i\}_{i \in [k]}$ .*

*Proof.* let  $L = \{\frac{\partial^2 f_i}{\partial x_a \partial x_b} \mid \frac{\partial^2 f_i}{\partial x_a \partial x_b} \neq 0\}_{i \in [k], a, b \in [n]}$ . Let  $P(x_1, \dots, x_n) = \prod_{g \in L} g(x_1, \dots, x_n)$ . By Lemma 3, each  $g \in L$  is a RO-ABP-polynomial. Hence by Lemma 1, for  $m = \lceil \log n \rceil + 1$ , the SV-generator  $(G_m^1, G_m^2, \dots, G_m^n)$ , satisfies that  $g(G_m^1, G_m^2, \dots, G_m^n) \neq 0$ , for all  $g \in L$ . So  $P(G_m^1, G_m^2, \dots, G_m^n) \neq 0$ .

Note that there are  $2m$  variables in  $P(G_m^1, \dots, G_m^n)$ , and the degree of every variable is bounded by  $kn^2 \cdot n^2 = kn^4$ . Thus by Lemma 5,  $\exists a \in V^{2m}, P(G_m^1(a), \dots, G_m^n(a)) \neq 0$ . Hence  $\mathcal{A}_k = G_n(V^{2m})$  is ensured to contain a nonzero of  $P$ . Any nonzero of  $P$  is a simultaneous nonzero of all  $g \in L$ . By Corollary 2,  $\mathcal{A}_k$  contains a simultaneous  $X$ -alignment for  $\{f_i\}_{i \in [k]}$ .  $\square$

## 7 A Hardness of Representation Theorem for RO-ABPs

The following theorem is an adaption of Theorem 6.1 in [2] to the notion of  $X$ -pre-alignment. One notable difference in the proof is that for the main case separation, we distinguish between whether there are 3rd-order partial derivatives vanishing or not (rather than 2nd-order partial as in [2]).

**Theorem 8.** *Assume  $|\mathbb{F}| > 3$ . Let  $P_n = \prod_{i \in [n]} x_i$ . If  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$  is a set of  $k$   $X$ -pre-aligned RO-ABP-polynomials for which  $P_n = \sum_{i \in [k]} f_i$ , then  $n < 7k$ .*

*Proof.* The proof proceeds by induction on  $k$ . For the base case  $k = 1$ , since  $f_1 = P_n$ , and  $f_1$  is  $X$ -pre-aligned, it must be that  $n \leq 2$ . Namely, if  $n > 2$ , then for  $x_i \in \text{Var}(P_n)$ , whatever distinct  $x_j, x_k \in X \setminus \{x_i\}$  we select,  $\frac{\partial^2 f_1}{\partial x_j \partial x_k} = x_i \cdot \prod_{x_r \in X \setminus \{x_i, x_j, x_k\}}$ . This cannot be of the form  $g \cdot (\beta x_i + \alpha)$  with  $g$  being an RO-ABP not depending on  $x_i$ , and  $\alpha = 0 \Rightarrow \beta = 0$ , as Definition 2 requires. Namely, since  $g$  does not depend on  $x_i$ , it must be that  $\beta \neq 0$ . Hence  $\alpha \neq 0$ , and thus  $g \cdot (\beta x_i + \alpha)$  is not homogeneous. Since  $x_i \cdot \prod_{x_r \in X \setminus \{x_i, x_j, x_k\}}$  is homogeneous, this is a contradiction.

Now assume  $k > 1$ . Suppose we can write  $P_n = \sum_{i \in [k]} f_i$ . For purpose of contradiction, assume that  $n \geq 7k$ . Hence  $n \geq 14$ .

**Case I:**  $\exists$  distinct  $p, q, r \in [n]$  and  $s \in [k]$ , such that  $\frac{\partial^3 f_s}{\partial x_p \partial x_q \partial x_r} \equiv 0$ .

Wlog. assume that  $p = n - 2, q = n - 1, r = n$  and  $s = k$ . Then  $\sum_{i \in [k-1]} \frac{\partial^3 f_i}{\partial x_{n-2} \partial x_{n-1} \partial x_n} = P_{n-3}$ .

By Lemma 8, all of the terms  $\frac{\partial^3 f_i}{\partial x_{n-2} \partial x_{n-1} \partial x_n}$  are  $(X \setminus \{x_{n-2}, x_{n-1}, x_n\})$ -pre-aligned. By induction, it must be that  $n - 3 < 5(k - 1)$ . Hence  $n < 5k - 2$ , which is a contradiction.

**Case II:**  $\nexists$  distinct  $p, q, r \in [n]$  and  $s \in [k]$ , such that  $\frac{\partial^3 f_s}{\partial x_p \partial x_q \partial x_r} \equiv 0$ .

We know  $\forall i, |\text{Var}(f_i)| \geq 3$ . Since  $f_i$  is  $X$ -pre-aligned, there exist distinct  $x_{j_i}, x_{k_i} \in X \setminus \{x_i\}$  such that  $\frac{\partial^2 f_i}{\partial x_{j_i} \partial x_{k_i}} = g_i \cdot (\beta_i x_n - \alpha_i)$ , where  $g_i$  is a RO-ABP-polynomial that does not depend on  $x_i$ , and  $\alpha_i = 0 \Rightarrow \beta_i = 0$ . Note that in this case,  $g_i \neq 0$ , since otherwise a second order partial vanishes. Hence both  $j_i$  and  $k_i$  are certainly not equal to  $x_n$ . It must be that  $\beta_i \neq 0$ , since otherwise  $\frac{\partial^3 f_i}{\partial x_{j_i} \partial x_{k_i} \partial x_n} \equiv 0$ . Hence also  $\alpha_i \neq 0$ .

**Claim 3.** Any  $g_i$  is  $(X \setminus \{x_{j_i}, x_{k_i}, x_n\})$ -pre-aligned.

*Proof.* Assume that  $|\text{Var}(g_i)| \geq 3$ , since otherwise the claim is trivial. Let  $h = g_i \cdot (\beta_i x_n - \alpha_i)$ . By Lemma 8,  $h$  is  $(X \setminus \{x_{j_i}, x_{k_i}\})$ -pre-aligned. Since  $\beta_i \neq 0$ , applying Corollary 1 yields that  $g_i$  is  $(X \setminus \{x_{j_i}, x_{k_i}, x_n\})$ -pre-aligned.  $\square$

Now, let  $A = \{\frac{\alpha_i}{\beta_i} : i \in [k]\}$ . Define for  $\gamma \in A$ ,  $E_\gamma = \{i \in [k] : \gamma = \frac{\alpha_i}{\beta_i}\}$  and  $B_\gamma = \{i \in [k] : \gamma \neq \frac{\alpha_i}{\beta_i} \text{ and } (f_i)|_{x_n=\gamma} \text{ is not } (X \setminus \{x_n\})\text{-pre-aligned}\}$ . Note that  $\sum_{\gamma \in A} |E_\gamma| = k$ . By Nearly Unique Nonalignment Lemma 11,  $\sum_{\gamma \in A} |B_\gamma| \leq 2k$ . Hence there exists  $\gamma_0 \in A$  such that  $|B_{\gamma_0}| \leq 2|E_{\gamma_0}|$ . Let  $I = E_{\gamma_0} \cup B_{\gamma_0}$ , and let  $J = \{j_i : i \in I\} \cup \{k_i : i \in I\}$ . We have that  $2 \leq |J| \leq 2|I| \leq 6|E_{\gamma_0}|$ . Observe that  $x_n \notin J$ . Define for any  $i$ ,  $f'_i = \partial_J f_i$ . We have the following three properties:

1. Each  $f'_i$  is an  $(X \setminus J)$ -pre-aligned RO-ABP-polynomial, due to Lemma 8.
2. For every  $i \in I$ ,  $f'_i = (\beta_i x_n - \alpha_i) h_i$ , where  $h_i$  is a RO-ABP-polynomial. Namely, since  $j_i, k_i \in J$ ,  $f'_i = \partial_{J \setminus \{j_i, k_i\}} [g_i (\beta_i x_n - \alpha_i)] = (\beta_i x_n - \alpha_i) \cdot \partial_{J \setminus \{j_i, k_i\}} g_i$ .
3. In the above, each  $h_i$  is an  $(X \setminus (J \cup \{x_n\}))$ -pre-aligned RO-ABP-polynomial. Namely, by Claim 3,  $g_i$  is  $(X \setminus \{x_{j_i}, x_{k_i}, x_n\})$ -pre-aligned. Hence, using Lemma 8, we get that  $h_i$  is an  $(X \setminus (J \cup \{x_n\}))$ -pre-aligned RO-ABP-polynomial.

For any  $i$ , define  $f''_i = (f'_i)|_{x_n=\gamma_0}$ . Then we have the following three properties:

1.  $\forall i \in E_{\gamma_0}$ ,  $f''_i \equiv 0$ .
2.  $\forall i \in B_{\gamma_0}$ ,  $f''_i = (\beta_i \gamma_0 - \alpha_i) h_i$ , so  $f''_i$  is an  $(X \setminus (J \cup \{x_n\}))$ -pre-aligned RO-ABP-polynomial, due to Proposition 4.

3. For every  $i \in [k] \setminus I$ ,  $(f_i)_{|x_n=\gamma_0}$  is  $X \setminus \{x_n\}$ -pre-aligned. Since  $n \notin J$ ,  $f_i'' = (f_i')_{|x_n=\gamma_0} = \partial_J[f_i]_{|x_n=\gamma_0}$ . So by Lemma 8,  $f_i''$  is an  $(X \setminus (J \cup \{x_n\}))$ -pre-aligned RO-ABP-polynomial.

Wlog. assume that  $J = \{\tilde{n} + 1, \tilde{n} + 2, \dots, n - 2, n - 1\}$ . Then  $|J| = n - 1 - \tilde{n}$ . Then  $\sum_{i \in [k]} f_i'' = (\partial_J P_n)_{|x_n=\gamma_0} = \gamma_0 \cdot P_{\tilde{n}}$ . Let  $\tilde{X} = \{x_1, \dots, x_{\tilde{n}}\}$ . We have found a representation of  $P_{\tilde{n}}$  as a sum of  $\tilde{k}$   $\tilde{X}$ -pre-aligned RO-ABP-polynomials, where  $7\tilde{k} \leq 7(k - |E_{\gamma_0}|) \leq n - 7|E_{\gamma_0}| = n - 1 - 6|E_{\gamma_0}| + 1 - |E_{\gamma_0}| \leq \tilde{n} + 1 - |E_{\gamma_0}| \leq \tilde{n}$ . This contradicts the induction hypothesis, and hence  $n < 7k$ .  $\square$

## 8 A Vanishing Theorem and the PIT Algorithms

The following theorem is analogous to Theorem 6.4 in [2].

**Theorem 9.** *Suppose  $|\mathbb{F}| > 3$ . Let  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$  be a set of  $k$   $X$ -aligned RO-ABPs. Let  $f = \sum_{i \in [k]} f_i$ . Then  $f \equiv 0 \iff f|_{\mathcal{W}_{7k}^n} \equiv 0$ .*

We need to argue only the “ $\Leftarrow$ ”-direction. Assume that  $f|_{\mathcal{W}_{7k}^n} \equiv 0$ .

We use induction on the number of variables  $n$ . The base case is when  $n < 7k$ . In this case it follows from Lemma 5 that  $f \equiv 0$ .

For the induction case assume  $n \geq 7k$ . We restrict one variable at a time. Consider a variable  $x_\ell$ , for  $\ell \in [n]$ . Consider a restriction of the polynomials  $f_i$ ’s and  $f$  to the subspace  $x_\ell = 0$ .

By condition 2 in the definition of *aligned*, each of the restricted polynomials  $f_i' = f_i|_{x_\ell=0}$  are  $(X \setminus \{x_\ell\})$ -aligned. Let  $f' = \sum_{i=1}^k f_i'$ . Clearly,  $f'|_{\mathcal{W}_{7k}^{n-1}} = f'|_{\mathcal{W}_{7k}^n} \equiv 0$ . Thus from the induction hypothesis,  $f' = f|_{x_\ell=0} \equiv 0$ , which implies that  $x_\ell$  divides  $f$ . Since  $\ell$  was arbitrarily chosen, this implies that  $P_n = \prod_{i=1}^k x_i$  divides  $f$ . But since  $f$  is multilinear, this gives  $f = c \cdot P_n$  where  $c$  is a constant and  $P_n = \prod_{i \in [n]} x_i$ .

Thus  $c \cdot P_n$  is the sum of  $k$  RO-ABPs which are also  $X$ -aligned (and therefore certainly  $X$ -pre-aligned). Since  $n \geq 7k$ , by Theorem 8, we can conclude that  $c = 0$ . Hence  $f \equiv 0$ .  $\square$

Now we are ready to give the identity testing algorithms for  $\Sigma_k$ -RO-ABP-polynomials given by  $\{f_i \in \mathbb{F}[X]\}_{i \in [k]}$ . The algorithm is simple. We use the fact that that  $\forall v \in \mathbb{F}^n$ ,  $f \equiv 0 \iff f(x_1 + v_1, x_2 + v_2, \dots, x_n + v_n) \equiv 0$ . Assuming that we have some common alignment  $v$  for  $\{f_i\}_{i \in [k]}$ , we know that each  $f_i(x_1 + v_1, x_2 + v_2, \dots, x_n + v_n)$  is  $X$ -aligned. In this case, Theorem 9 is applicable, and it suffices to test if the polynomial evaluates to zero on the set  $\mathcal{W}_{7k}^n$ . Based on the three approaches to get a common alignment, the algorithms are as follows:

1. (*Non-black-box setting*) By Corollary 3, we obtain a simultaneous alignment in time  $O(k^2 n^7 s)$ . Then it takes  $n^{O(k)}$  to test all points in  $\mathcal{W}_{7k}^n$ , so the running-time is  $O(k^2 n^7 s) + n^{O(k)}$ . This proves Theorem 4. In this case we need  $|\mathbb{F}| > kn^2$ .
2. (*Semi-black-box setting*) By Corollary 4, we obtain a simultaneous alignment in time  $k^2 n^{O(\log n)}$ . Then it takes  $n^{O(k)}$  to test all points in  $\mathcal{W}_{7k}^n$ , so the running-time is  $k^2 n^{O(\log n)} + n^{O(k)}$ . This proves Theorem 5. In this case we need  $|\mathbb{F}| > kn^2$ .
3. (*Black-box setting*) In this case we only have black-box access to  $f = \sum_{i \in [k]} f_i$ . Let  $f_v(x_1, \dots, x_n) = f(x_1 + v_1, \dots, x_n + v_n)$ . Then it is easy to see that  $f \equiv 0 \iff \forall v \in \mathcal{A}_k, f_v|_{\mathcal{W}_{7k}^n} \equiv 0$ . In this case the running-time is  $n^{O(\log n + k)}$ . This proves Theorem 2. In this case we need  $|\mathbb{F}| > kn^4$ .

## References

- [1] A. Shpilka and I. Volkovich. Read-once polynomial identity testing. In *Proceedings of the 40th Annual STOC*, pages 507–516, 2008.
- [2] A. Shpilka and I. Volkovich. Improved polynomial identity testing of read-once formulas. In *Approximation, Randomization and Combinatorial Optimization. Algorithms and Techniques, volume 5687 of LNCS*, pages 700–713, 2009.
- [3] M. Agrawal. Proving lower bounds via pseudo-random generators. In *Proc. 25th Annual Conference on Foundations of Software Technology and Theoretical Computer Science*, pages 92–105, 2005.
- [4] N. Saxena. Progress of polynomial identity testing. Technical Report ECCC TR09-101, Electronic Colloquium in Computational Complexity, 2009.
- [5] J.T. Schwartz. Fast probabilistic algorithms for polynomial identities. *J. Assn. Comp. Mach.*, 27:701–717, 1980.
- [6] R. Zippel. Probabilistic algorithms for sparse polynomials. In *Proceedings of the International Symposium on Symbolic and Algebraic Manipulation (EUROSAM '79)*, volume 72 of *Lect. Notes in Comp. Sci.*, pages 216–226. Springer Verlag, 1979.
- [7] V. Kabanets and R. Impagliazzo. Derandomizing polynomial identity testing means proving circuit lower bounds. *Computational Complexity*, 13(1–2):1–44, 2004.
- [8] M. Ben-Or and P. Tiwari. A deterministic algorithm for sparse multivariate polynomial interpolation. In *Proc. 20th Annual ACM Symposium on the Theory of Computing*, pages 301–309. ACM, 1988.
- [9] A.R. Klivans and D.A. Spielman. Randomness efficient identity testing of multivariate polynomials. In *Proc. 33rd Annual ACM Symposium on the Theory of Computing*, pages 216–223, 2001.
- [10] R. Lipton and N. Vishnoi. Deterministic identity testing for multivariate polynomials. In *Proceedings of the fourteenth annual ACM-SIAM symposium on Discrete algorithms (SODA 2003)*, pages 756–760, 2003.
- [11] Z. Dvir and A. Shpilka. Locally decodable codes with two queries and polynomial identity testing for depth 3 circuits. *SIAM J. Comput.*, 36(5):1404–1434, 2006.
- [12] N. Kayal and N. Saxena. Polynomial identity testing for depth 3 circuits. *Computational Complexity*, 16(2):115–138, 2007.
- [13] V. Arvind and P. Mukhopadhyay. The monomial ideal membership problem and polynomial identity testing. In *Proceedings of the 18th International Symposium on Algorithms and Computation (ISAAC 2007)*, volume 4835 of *Lecture Notes in Computer Science*, pages 800–811. Springer, 2007.

- [14] Z.S. Karnin and A. Shpilka. Deterministic black box polynomial identity testing of depth-3 arithmetic circuits with bounded top fan-in. In *Proc. 23rd Annual IEEE Conference on Computational Complexity*, pages 280–291, 2008.
- [15] N. Kayal and S. Saraf. Black box polynomial identity testing of depth-3 circuits. In *Proc. 49th Annual IEEE Symposium on Foundations of Computer Science*, 2009.
- [16] Z.S. Karnin, P. Mukhpadhyay, A. Shpilka, and Ilya Volkovich. Deterministic identity testing of depth 4 multilinear circuits with bounded top fan-in. Technical Report TR09–116, Electronic Colloquium on Computational Complexity (ECCC), November 2009.
- [17] R. Raz and A. Shpilka. Deterministic polynomial identity testing in non commutative models. *Computational Complexity*, 14(1):1–19, 2005.
- [18] M. Agrawal and V. Vinay. Arithmetic circuits: A chasm at depth four. In *Proc. 49th Annual IEEE Symposium on Foundations of Computer Science*, pages 67–75, 2008.
- [19] L. Valiant. Completeness classes in algebra. Technical Report CSR-40-79, Dept. of Computer Science, University of Edinburgh, April 1979.
- [20] L. Lovász. On determinants, matching, and random algorithms. In *FCT’79: Fundamentals of Computation Theory*, pages 565–574, 1979.
- [21] K. Mulmuley, U. Vazirani, and V. Vazirani. Matching is as easy as matrix inversion. *Combinatorica*, 7:105–113, 1987.
- [22] E. Allender, K. Reinhardt, and S. Zhou. Isolation, matching and counting uniform and nonuniform upper bounds. *J. Comput. Syst. Sci.*, 59(2):164–181, 1999.
- [23] S. Datta, R. Kulkarni, and S. Roy. Deterministically isolating a perfect matching in bipartite planar graphs. In *Proc. 25th Annual Symposium on Theoretical Aspects of Computer Science*, volume 08001 of *Leibniz Int. Proc. in Informatics*, pages 229–240, 2008.
- [24] M. Jansen. Weakening assumptions for deterministic subexponential time non-singular matrix completion, 2010. To Appear, 27th International Symposium on Theoretical Aspects of Computer Science (STACS 2010).
- [25] N. Alon. Combinatorial nullstellensatz. *Combinatorics, Probability and Computing*, 8(1–2):7–29, 1999.

## A Figure 3

Figure 3 shows an RO-ABP computing  $x_1x_2 + x_2x_3 + x_{n-1}x_n$ , when  $n$  is even. The case when  $n$  is odd is dealt with similarly. Unlabeled edges are labeled with 1.

## B Example : RO-ABPs Are Not Universal

**Proposition 6.** *The degree-2 elementary symmetric polynomial  $e_n(x_1, x_2, \dots, x_n) = \prod_{1 \leq i < j \leq n} x_i x_j$ ,  $n \geq 3$  can not be computed by a RO-ABP.*




Figure 3: A RO-ABP computing  $x_1x_2 + x_2x_3 + \dots + x_{2n-1}x_{2n}$ .

*Proof.* For the purpose of contradiction, suppose that some RO-ABP  $A$  computes  $e_n$ . For any  $x_i$  denote the edge it labels by  $g_i = (s_i, t_i)$ . We can define an ordering  $<$  among  $g_i$ 's, by taking  $g_i < g_j$  if and only if the polynomial computed by the subprogram  $A(t_i, s_j)$  has a nonzero constant term. Due to the fact that  $A$  is a DAG, we have for any  $i, j$ , if  $x_i < x_j$ , then not  $x_j < x_i$ .

The fact that for every  $(i, j)$  pair,  $x_i x_j$  appears as a term in  $e_n$  implies that for any  $i \neq j$ , we have one of  $x_i < x_j$  or  $x_j < x_i$ . Incidentally, note this implies the ordering is transitive. Namely, if  $x_i < x_j$  and  $x_j < x_k$ , then  $s_j$  must be reachable from  $t_i$ , and  $s_k$  must be reachable from  $t_j$  in  $A$ , but then  $s_i$  can not be reachable from  $t_k$ . Hence not  $x_k < x_j$ , which implies  $x_j < x_k$ .

In any case, observe there is a permutation  $\phi : [n] \rightarrow [n]$  for which  $x_{\phi(1)} < x_{\phi(2)} < \dots < x_{\phi(n)}$ . This implies that  $\prod_{i \in [n]} x_i$  appears as a term in the polynomial computed by  $A$ , which is a contradiction.  $\square$

## C Non-Black-Box Testing a Single RO-ABP

Consider a RO-ABP  $A$ . Denote the source and sink of  $A$  by  $s$  and  $t$ , respectively. Suppose that  $x_i$  labels the edge  $(s_i, t_i)$ . Wlog. assume that the order of variable layers in  $A$  is  $x_1, x_2, \dots, x_n$ . We have the following easy proposition:

**Proposition 7.** *Suppose  $1 \leq i_1 < i_2 < \dots < i_k \leq n$ . For a RO-ABP  $A$ ,  $x_{i_1} x_{i_2} \dots x_{i_k}$  appears in  $\hat{A}$  if and only if the constant terms in  $\hat{A}(s, s_{i_1})$ ,  $\hat{A}(t_{i_m}, s_{i_{m+1}})$ , for all  $m \in [k-1]$ , and  $\hat{A}(t_k, t)$  are not zero.*

We build a directed graph  $G_A = (V, E)$  for RO-ABP  $A$  with vertex set  $V = \{s, t, x_1, x_2, \dots, x_n\}$ . Edges are given as follows:

1.  $(s, x_i)$ , if the constant term in  $\hat{A}(s, s_i)$  is nonzero.
2.  $(x_i, t)$ , if the constant term in  $\hat{A}(t_i, t)$  is nonzero.
3.  $(x_i, x_j)$ ,  $i < j$ , if the constant term in  $\hat{A}(t_i, s_j)$  is nonzero.

We have the following corollary of Proposition 7:

**Corollary 5.**  $\hat{A}(x_1, \dots, x_n) \equiv 0$  if and only if  $t$  is not reachable from  $s$  in  $G_A$ .

The algorithm for testing  $A$  is to construct  $G_A$  and to test connectivity. This can be done in time  $O(n^2 s)$ , where  $s$  bounds the size of  $A$ .